

2019-04-16

Assessing the impact of introduced infrastructure at sea with cameras: A case study for spatial scale, time and statistical power

Bicknell, AWJ

<http://hdl.handle.net/10026.1/13760>

10.1016/j.marenvres.2019.04.007

Marine Environmental Research

Elsevier

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Assessing the impact of introduced infrastructure at sea with cameras: a case study for spatial scale, time and statistical power

Anthony W. J. Bicknell, (Present) Environmental Biology, University of Exeter, Hatherly Building, Prince of Wales Road, Exeter, EX4 4PS, UK, Environment and Sustainability Institute, University of Exeter, Penryn, Cornwall TR10 9FE, UK, and Centre for Ecology and Conservation, University of Exeter, Penryn, Cornwall TR10 9FE, UK a.bicknell@exeter.ac.uk (corresponding author)

Emma V. Sheehan, School of Biological and Marine Sciences, University of Plymouth, Plymouth, Devon PL4 8AA, UK emma.sheehan@plymouth.ac.uk

Brendan J. Godley, Centre for Ecology and Conservation, University of Exeter, Penryn, Cornwall TR10 9FE, UK B.J.Godley@exeter.ac.uk

Philip D. Doherty, Centre for Ecology and Conservation, University of Exeter, Penryn, Cornwall TR10 9FE, UK P.Doherty@exeter.ac.uk

Matthew J. Witt, (Present) Environmental Biology, University of Exeter, Hatherly Building, Prince of Wales Road, Exeter, EX4 4PS, UK, Environment and Sustainability Institute, University of Exeter, Penryn, Cornwall TR10 9FE, UK, and Centre for Ecology and Conservation, University of Exeter, Penryn, Cornwall TR10 9FE, UK M.J.Witt@exeter.ac.uk

Abstract

Detecting the effects of introduced artificial structures on the marine environment relies upon research and monitoring programs that can provide baseline data and the necessary statistical power to detect biological and/or ecological change over relevant spatial and temporal scales. Here we report on, and assess the use of, Baited Remote Underwater Video (BRUV) systems as a technique to monitor diversity, abundance and assemblage composition data to evaluate the effects of marine renewable energy infrastructure on mobile epi-benthic species. The results from our five-year study at a wave energy development facility demonstrate how annual natural variation (time) and survey design (spatial scale and power) are important factors in the ability to robustly detect change in common ecological metrics of benthic and bentho-pelagic ecosystems of the northeast Atlantic. BRUV systems demonstrate their capacity for use in temperate, high energy marine environments, but also how weather, logistical and technical issues require increased sampling effort to ensure statistical power to detect relevant change is achieved. These factors require consideration within environmental impact assessments if such survey methods are to identify and contribute towards the management of potential positive or negative effects on benthic systems.

Keywords

Marine monitoring, human impact, renewable energy, power analysis, BRUV

Introduction

The marine coastal environment provides a major, and disproportionate, contribution to global ecosystem services (Costanza et al., 1997; Drakou et al., 2017; Lique et al., 2013). Through human activity (e.g. commercial fishing, shipping, resource extraction, aquaculture, dredging) it has, however, become the most impacted region of our global seas (Halpern et al., 2008). A growing coastal population is expected to increase human pressure on coastal regions with further potential adverse effects on natural systems (Heery et al., 2017; Millennium Ecosystem Assessment, 2005; Neumann et al., 2015).

Coastal benthic habitat has been particularly altered by destructive fishing practices (Eigaard et al., 2015; Hiddink et al., 2017) and the introduction of artificial infrastructure (Bulleri and Chapman, 2010). The loss or disturbance of benthic habitats is concerning as they play a pivotal role in the provision and support of key ecosystem services, e.g. food provision, nutrient cycling, reproduction/nursery areas, water quality, biodiversity maintenance (Galparsoro et al., 2014). The expanding marine renewable energy sector has led offshore wind farms to become a prominent part of coastal and shelf waters of multiple countries (GWEC, 2016; WindEurope, 2017). These installations have direct impact on benthic habitats (i.e. monopile drilling and foundations), and subsequent direct or indirect effects on the associated epi-benthic faunal communities (Bailey et al., 2014; Gill, 2005; Inger et al., 2009; Pearce et al., 2014; Stenberg et al., 2015). The promise of wave and tidal energy conversion is still to be fully realised, but has the potential to further modify benthic habitats around our coastlines (Langhamer and Wilhelmsson, 2009; Witt et al., 2012). Unfortunately, the monitoring programs and environmental impact assessments (EIA) that have considered interactions between marine renewable energy installations (MREIs) and benthic habitats or species have, so far, lacked the necessary baseline characterisation or survey rigour (spatially or temporally) to robustly assess impact (positive or negative) (Boehlert and Gill, 2010; Fox

et al., 2018; Wilding et al., 2017), and led authors to term this situation as ‘data rich, information poor (DRIP)’ (Fox et al., 2018; Ward et al., 1986; Wilding et al., 2017).

The high energy environment needed for marine energy convertors to operate means they are well suited to mid to high latitude seas where these conditions regularly occur. The required strong winds, wave action and/or large tides provide physical challenges to access development sites and conduct surveys to assess impact on benthic communities, which are in addition to the inherent difficulties of underwater research. Remote camera imagery is a technique that can overcome some of these challenges, and has already proven a valuable tool for studying the impact of human activities on the marine environment (Bicknell et al., 2016; Mallet and Pelletier, 2014; Sheehan et al., 2014). Baited Remote Underwater Video systems (known as ‘BRUV’) are a method that uses either one (mono) or two (stereo) cameras to film the area surrounding a bait attractant held a short distance from a video camera and close to the seabed (also modified for mid-water; Heagney et al., 2007). The technique has been used extensively in the southern-hemisphere (tropics to temperate) to evaluate changes in demersal fish populations (e.g. Denny et al., 2004; Malcolm et al., 2007; Watson et al., 2009; Watson et al., 2007), and has demonstrated its value in sampling fishes and invertebrates in high latitude turbid coastal waters in the northern-hemisphere (Elliott et al., 2017; Unsworth et al., 2014). Recently, the method has also shown its application in assessing the mobile epibenthic fauna at an offshore wind farm in the Irish Sea (Griffin et al., 2016), but as yet not in multi- season or multi- year impact studies. The method has bias (e.g. differentially attracting carnivores or omnivores, bait type and plume effects, restricted view, light attraction/repulsion), as do most survey methods, and these have been investigated and detailed elsewhere (Dorman et al., 2012; Harvey et al., 2018; Harvey et al., 2012; Harvey et al., 2007; Stobart et al., 2007). However, it has advantages on many traditional methods, such as being non-destructive, having no or limited observer bias, allowing re-analysis or review

of video (data) and is unrestricted by depth (cost-dependent) (Cappo et al., 2004; Lowry et al., 2012; Whitmarsh et al., 2016; Zintzen et al., 2012). When BRUVs are used to provide an estimate of species abundance, a number of metrics have been considered (Stobart et al., 2015), but in the vast majority of cases (81% of reviewed studies, Whitmarsh et al., 2016) N_{max} (or MaxN) is used. This represents the maximum number of a particular species seen in any one video frame across the duration of the video footage. It is a useful metric to assess the relative abundance of species and considered a conservative estimate as there may be uncounted individuals around the BRUV that did not enter the field of view (Whitmarsh et al., 2016).

For monitoring programs to effectively assess environmental impact they require baseline data that characterise the natural spatial and temporal variability of the focal system or component (Judd, 2012). The challenges are then to detect the potential effect of the introduced impact from the natural ‘background noise’ (variability) (Osenberg et al., 1994), and recognize whether any detected change is biologically, ecologically or functionally meaningful (Wilding et al., 2017). In highly variable marine systems it is particularly important to determine the level of sampling effort required to gather robust baseline data and provide statistical power to detect a given degree of change (Franco et al., 2015; Osenberg et al., 1994). Ideally, prior or pilot data would be available to provide knowledge on variability within the system and enable a power analysis to be conducted, but these data are rarely available or analyses conducted (Franco et al., 2015; Maclean et al., 2014). Surveys will often be based on applying fundamental statistical principles to the design (Box 1), balanced with time, costs and logistical or methodological constraints. Many environmental impact studies focus on site characterisation during one or two years as opposed to deploying bespoke survey strategies designed to identify putative effects with anticipated levels of change in environmental receptor groups.

Effects of design and data on statistical power

The ability to detect patterns/change reduces as variability in the parameter being measured increases

↑variance = ↓power

Parameter estimates become more precise with larger samples, thus differences between estimates are easier to detect amongst the 'noise'

↑sampling = ↑estimate precision

↑estimate precision = ↑power

As the effect of the impact increases the more likely it is to be detected. i.e. a 40% change in a parameter estimate is more likely to be detected than 20% change

↑effect size = ↑power

(Underwood and Chapman, 2003)

Here, we present a case study of BRUV use over five years in boreal latitude coastal waters of the northeast Atlantic to provide baseline characterisation data and impact assessment on mobile epi-benthic species at a MREI. We use these data to investigate the power to detect change in conventional ecological metrics (species richness, abundance, and assemblage composition), how well a survey design and sampling effort performs given there was little prior ecological knowledge of the site, and whether any effects on the mobile epi-benthic community could be detected. We consider the appropriateness of this technique for long-term impact monitoring at MREI, and, more generally, discuss the results in relation to future EIAs of mobile epi-benthic communities.

Materials and Methods

Study location

Baited remote underwater video (BRUV) surveys took place off the north coast of Cornwall (UK) between 2011 and 2015. The study was located within and adjacent to a MREI development zone (Wave Hub) and the associated seabed cable (Fig. 1a). Study zones ranged between 3 and 10.5 nautical miles (5.6 and 19.5 km) offshore in water depths of 20 to 53 m

(at Lowest Astronomical Tide). The seabed cable and marker buoys (6 in total) for the development zone were installed in autumn 2010. The cable was buried when on sand (near shore) and covered with 80,000 tonnes of rock (creating a berm of 0.3 metres minimum height) and concrete matressing every 120 metres when laid on hard substrate (deeper water). Access within 500 metres of the electrical seabed hub (plug) was prohibited, but access was permitted elsewhere. The presence of marker buoys has prevented commercial mid-water or bottom trawling for fish. The only commercial fishing that has taken place in the development zone or close to the seabed cable since installation is crustacean potting. There were no connected deployments of operating marine energy devices during the period of the study.




Fig. 1. Study zones and sampling locations (green filled circles) for BRUV surveys (a), BRUV housing, frame and LED light (b), and example habitat types from the video footage (c) 1 = rocky reef, 2 = large (course) sediment, 3 = medium (mixed) sediment, 4 = fine sediment.

Sampling equipment

Each BRUV consisted of an aluminium frame, wide-angle lens housing and white light LED lighting system (Fig. 1b). An aluminium pole was attached to each BRUV to support bait (fixed 1 metre from lens); lead weights (45 kg) were fastened to the frame for deployment and stability over a wide range of tidal conditions (0.02 to 0.53 m^{-s}). Panasonic HDC-SD60 and HDC-SD80 camcorders were used to gather video data. A temperature depth recorder (RBR, Nova Scotia, Canada) was attached to one BRUV in each sampling location during a sampling campaign. Local small commercial leisure or fishing vessels were used for BRUV deployment.

Survey and design

BRUV surveys were scheduled twice per year, in spring and autumn, commencing in autumn 2011 until 2015, and each survey campaign took three days to complete. The sampling design consisted of: two study *zones* (Wave Hub = WHSZ and Cable Route = CRSZ); three *areas* (1 x Wave Hub, 2 x Cable Route); each area comprised 8 *locations* (four **project** and four **reference** 1 km apart; Fig 1a). Three replicate BRUVs were deployed between 105-180 m (mean ~130 m) apart in each location (Fig 2). **Project** locations were either inside the Wave Hub exclusion zone or on/next to the seabed cable infrastructure (Fig 1a). *Treatment* will be used as the term to describe the comparison between **project** and **reference** locations in the subsequent analyses and models. BRUVs were deployed for up to 60 minutes during daylight hours. Bait used was a single Atlantic mackerel *Scomber scombrus* for each deployment, cut into three piece and held in a net bag (~100 g).

In order to investigate the BRUVs greatest distance of attraction (in metres) for teleosts, we calculated the ‘effective range of attraction’ (AR) (formulised in Cappo et al., 2004; see Appendix S3) for increasing soak time (i.e. time cameras are in the water, or video footage analysed from the start). The average seabed current speeds (data from POLPRED seabed

CS20 models, NOC; <http://www.pol.ac.uk/>) during BRUV deployments (0.23 m s^{-1}), and a maximum fish (endurance) swimming speed of 0.6 m s^{-1} (~200-300 mm fish length) were used as the AR parameters (V_c = current speed, V_f = maximum fish speed).




Fig. 2. Nested design schematic for baited remote underwater video surveys.

Image analysis

Gathered video datasets were analysed (using a large monitor) to quantify species observed, the number of mobile epi-benthic species (richness; S), the maximum number of individuals of each species observed at the same time (N_{max}) and the time of each increment in N_{max} (recorded in excel spreadsheets). The use of N_{max} as an estimator of relative abundance has been assessed (Cappo et al., 2003; Ellis and DeMartini, 1995; Priede et al., 1994; Willis and Babcock, 2000), and is considered a conservative estimate of abundance especially when species occur at high density. Each video dataset was assigned a habitat type (1-rocky reef, 2-large sediment [small boulders], 3-medium sediment, 4-fine sediment; see Appendix S2 for habitat assignment details), visibility (good [can see beyond end of pole] or poor [could not see beyond end of pole]) and camera frame position (vertical or horizontal).

Data analysis

Data on all mobile species were used in the following S analyses but only teleosts were used in the N_{max} analyses. To remove the influence of extremely high abundance values, the teleost N_{max} data were trimmed at the 95th percentile (eliminating 3 data points). Extreme abundance values were related to large fish shoals that were rare and not informative for the purpose of determining subtle and consistent change in species abundance.

Species accumulation

To examine the effect of BRUV deployment time on S and N_{max} , species and abundance accumulation curves were created for the Wave Hub and Cable Route (project and reference location data pooled). To investigate the potential effect of habitat, species and abundance accumulation curves were created with all data pooled for habitat types; rocky reef and large sediment (termed RRLS), and medium (gravel) and fine sediment (sand) (termed GS). These curves were used to determine an optimum balance between the number of available comparative video datasets and their recording duration (minutes) to be included in subsequent analyses. All conducted in R version 3.4.4 (R Development Core Team, 2011)

Statistical power, effect and sample size

Power analyses were conducted using the SIMR package (Green and MacLeod, 2016) in R version 3.4.4 (R Development Core Team, 2011) to investigate the relationships between the number of samples (sample size), the size of change (effect size) and the probability to detect change (power) in S or N_{max} .

Generalised linear mixed effect models were fitted to data (pooled data with years combined) collected in spring at the WHSZ and CRSZ (separate models) using the lme4 package (Bates et al., 2015). Only spring data were used in the models to remove seasonal effects (evidenced in PERMANOVA models; Table 3a & 3b), which may increase variability in the data and

influence the power to detect change. (Step 1) The Poisson error models contained *year* and *treatment* as fixed effects and *treatment* within *location* as a nested random effect. The effect parameters from these models were used to build new simulation models with the same structure (i.e. effects and error) but replicated a complete annual survey design for each study zone (i.e. no loss of samples). (Step 2) Monte Carlo simulation was then utilised to generate values for the response variable (S or N_{max}) of each model (1000 runs & seed = 1234). (Step 3) Power curves were generated for a range of effect sizes to explore the trade-off between sample size and power. (Step 4). To examine inter-annual variation the steps detailed above (1-3) were repeated for separate years (WHSZ = 4 years, CRSZ = 3 years) for effect sizes where the pooled data model reached or exceeded 0.8 power. All analyses were conducted using an (α) <0.05 threshold significance level, and results related to 0.8 power, the commonly accepted level of confidence.

Species richness, abundance and assemblage composition

Permutational multivariate and univariate mixed effect models (PERMANOVA+) were used in the software package PRIMERv6 (Anderson, 2001; Clarke and Warwick, 2001) to test potential effects of the Wave Hub exclusion zone and the cable rock armouring (Cable Route) on assemblage composition (a community structure measure incorporating both diversity and abundance), species richness (S) and abundance (N_{max}).

Between study zones

Models using the complete dataset were run to compare assemblage composition and S between study zones (WHSZ & CRSZ). Factor *zone* was fixed, with *year*, *season*, *location*, and *treatment* nested in *location* as random.

Within study zones

For each study zone (WHSZ & CRSZ), models for S and assemblage composition were performed on the complete species dataset, and all three response variables for two defined species guilds; teleost and crustacean. The factors *year* and *treatment* were fixed, and *season*, *habitat*, *zone* (Cable Route only; 2 zones) and *treatment* nested in *location* were random. The factor *year* had four and five levels for the CRSZ and the WHSZ respectively. The factor *treatment* had two levels (WHSZ or CRSZ project, and reference). Depth and current speed (tide indicator) were considered as environmental co-variates in initial models for each study zone but were removed when found not to influence models.

Species N_{max} models were performed for four indicator taxa, family and species: elasmobranchs, echinoderms, and *Pollachius spp.* (*Pollack* & *Saithe*) and *Cancer pagurus* (*edible crab*) (Jackson et al., 2009). The same model structure was used as described above.

Between habitats

Models were also run using the complete dataset to test whether S , N_{max} and assemblage composition differed with habitat type. Due to the low number of sampling events on fine sediment habitats (n=9), making the model unbalanced, the final model contained only 3 levels (rocky reef [n=84], large [n=47] and medium [n=58] sediment) for the fixed factor *habitat*, with *year* as random.

Prior to calculation of the Bray–Curtis (Bray and Curtis, 1957) similarity index, multivariate data (assemblage composition) were dispersion weighted and square root transformed to down weight taxa with erratic abundances and/or high abundances (Clarke et al., 2006a). As joint species absences were important to consider between treatments, data were ‘zero-adjusted by adding a dummy value of 1 (Clarke et al., 2006b). Without the dummy value, Bray-Curtis would not consider samples similarly devoid of species as similar. Euclidean

distance indices were calculated for univariate data (S and N_{max}) that were $\text{Log}(x + 1)$ transformed (Anderson and Millar, 2004). Each term in the analyses used 9999 permutations of the appropriate units. Significant interactions of fixed terms were tested using PERMANOVA pairwise tests. Assemblage composition was visualised using nonmetric Multi-Dimensional Scaling (nMDS).

Results

Sampling and image quality

The sampling regime was designed to gather data in both spring and autumn of each year (2011-2015; full survey design = 720 deployments) to examine the potential effects of season, but due to poor weather conditions in optimal tidal survey periods, the collection of data in both seasons was only possible in 2013. Autumn sampling took place in 2011 and 2013; spring sampling took place in 2012, 2013, 2014 and 2015. Due to time and budgetary constraints, no surveys could be completed in 2015 at the CRSZ, and the original survey extent was reduced by two locations (most southerly project and reference locations in each area) in 2013-2015 at the WHSZ and 2012-2014 for the CRSZ. The reduced survey effort owing to weather, time and money resulted in a potential maximum of 312 BRUV deployments, a considerable reduction (57%) from the original sampling design.

Further adverse weather during survey activity caused 15 (5%) deployments to be cancelled. Across study areas and years, 297 BRUV deployments were successfully undertaken, of which 38 (12.8%) failed due to technical reasons (e.g. battery or camera failure). Of the remaining 259 (83% of total); 141 video datasets were 60 minutes or longer (45% of total), 199 video datasets were 45 minutes or longer (63% of total), and 247 were 30 minutes or longer (79% of total). Once filtered for visibility (good visibility with an unobstructed view of the seabed) the number of available videos for analysis were 116 with 60 minutes or more

300 (37% of total), 161 datasets with 45 minutes or more (51% of total), and 198 datasets with 30
301 minutes or more (63% of total) (Fig. S2).

302 The BRUVs effective range of attraction (AR) for a teleost of ~200-300 mm in length was
303 ~127 m for 30 minutes soak time, ~287 m for 45 minutes and ~510 m for 60 minutes for our
304 study (Fig. S3). This would indicate the replicate BRUV deployments in our study (mean
305 ~130 m apart) may not be independent (i.e. the same individual could attend and be recorded
306 on more than one camera) if more than 30 minutes video data were used. The AR calculation
307 does not consider current or plume direction, which could influence the range of attraction
308 shape (e.g. not cylindrical around the location, but elongated (ellipsoid/triangular) in the
309 direction of current). This in turn could affect the ability of animals to detect bait from a
310 second location (reduce the AR estimate in that direction) while visiting the first, if the bait
311 plume is directed away from the first. The BRUV replicates in our study were placed in a
312 linear manner across the prevalent NE↔SW tide, mainly N↔S and NW↔SE (Fig. 1). The
313 current direction and speed data for each BRUV deployment (Fig. S4) indicates the prevalent
314 seabed currents would advect bait plume away from other replicate locations rather than
315 towards them, suggesting it would be more difficult for an animal to detect bait from a
316 second location and subsequently move to it. When 30 minutes video data were analysed for
317 spatial auto-correlation (Moran's I) using teleost and elasmobranch abundance for each study
318 zone, year and season, the only significant auto-correlation was found for teleost in 2012 (2
319 out of 17 tests; Table S1). When the data for 2012 were investigated further, the effect was
320 likely caused by large aggregations of *Trisopterus minutus* counted on all three replicate
321 BRUVs at locations in each study zone in this year. This is a small (~100-200 mm) benthic
322 species, so their ability to attend two or three BRUVs (linearly ~130 or 260 m apart) would
323 be unlikely given the AR and current data, and the clustering (auto-correlation) would seem
324 more likely a representation of high abundance at all these locations during that time. Given

these analyses it was considered that species and abundance data gathered from 30 minutes video data would be independent for teleost and the (relatively) small elasmobranch species in our study.

Species accumulation

The maximum number of species identified during video datasets lasting 60 minutes was 12. The mean number of species occurring on RRLS habitats was 6.8, while for GS it was 5.3 (Fig. 3a & c). Mean N_{max} was greater for RRLS habitats (maximum of 132 individuals) compared to GS habitats (maximum 38 individuals) (Fig. 3b & d). In both habitats, 75% or more of the total S and N_{max} (at 60 minutes) was achieved after 30 minutes, with the greatest increase in number of species and abundance (rate of change) occurring in the first 20 minutes of video datasets (Fig S1).




Fig. 3. Mean species (a & c) and relative abundance (N_{max}) (b & d) accumulation curves from 60 minute BRUV footage for rocky reef/large sediment habitat (a & b) and medium/fine sediment habitats (c & d). Grey shading = 95% confidence interval. Dashed lines = 30, 45 and 60 minutes.

Optimal data

Data from the first 30 minutes of BRUV footage were considered optimal for use in further analyses as this soak period balanced sample size, coverage (study locations and years), effective range of attraction, with available species and abundance information (accumulation curves).

Species diversity and taxonomic composition

A total of 67 species from 46 families and 6 phyla were observed on the BRUV footage across the two study zones for the survey period. This equated to 5,440 individual animals, the vast majority of which were fishes (teleost ~73%). The WHSZ was more speciose (49 species) compared to the CRSZ (42 species), but total abundance was higher in the latter with >3400 individuals (Table 1).

Statistical power, effect and sample size

Models with pooled spring data (years combined) revealed the minimum change that could be detected with 0.8 (or more) power was ~10% in richness for both WHSZ and CRSZ, which would require ≥ 200 samples (~100 within each project and reference). The samples required to detect 20% change in S reduce to ~40-50 (~20-25 within each project and reference) for both WHSZ and CRSZ (Fig 4a & 4c sub plots). For N_{max} , the minimum change detectable was ~50% at the WHSZ (Fig 4b sub plot), which required ≥ 100 samples (~50 within each project and reference). The 0.8 power threshold was not reached for any level of

363 change in N_{max} using pooled spring data from the CRSZ, and the power to detect 50% change
364 did not exceed 0.4, no matter the sample size (Fig 4d sub plot).

365 **Table 1** Number of species and individuals by taxonomic phylum on BRUV footage at the Wave Hub and Cable Route study zones. The
 366 percentage of total species and individuals is shown in italics.

Phylum	Wave Hub Study Zone				Cable Route Study Zone			
	No. of species	<i>% of total species</i>	No. of individuals	<i>% of total individuals</i>	No. of species	<i>% of total species</i>	No. of individuals	<i>% of total individuals</i>
Arthropod	1	<i>2.00</i>	2	<i>0.10</i>	0	<i>0.00</i>	0	<i>0.00</i>
Crustacean	13	<i>26.53</i>	331	<i>16.62</i>	8	<i>19.05</i>	165	<i>4.79</i>
Echinoderm	6	<i>12.25</i>	261	<i>13.10</i>	6	<i>14.29</i>	390	<i>11.31</i>
Elasmobranch	3	<i>6.12</i>	162	<i>8.13</i>	2	<i>4.76</i>	152	<i>4.41</i>
Mollusc	5	<i>10.20</i>	18	<i>0.90</i>	0	<i>0.00</i>	0	<i>0.00</i>
Teleost	21	<i>42.86</i>	1218	<i>61.14</i>	26	<i>61.90</i>	2741	<i>79.50</i>
Total	49		1992		42		3448	

367

368




Fig. 4. Minimum effect size SIMR model outputs ($\pm 95\%$ CI; <0.05 significance) for pooled spring species richness (S) and relative teleost abundance (N_{max}) data (inserted plots) separated by year. (a) = 10% change in S at WHSZ, (b) = 10% change in S at CRSZ, (c) = 50% change in N_{max} at WHSZ, and (d) = 50% change in N_{max} at CRSZ. Green dashed line = 2012, blue dashed line = 2013, purple dashed line = 2014, and orange dashed line = 2015. Dot-dash line on all plots indicates 0.8 power.

There was considerable variation among years when data were analysed separately for 20% (S) and 50% (N_{max}) effect size models (detailed above; Fig. 4). The sample size required to detect a 20% change in S at WHSZ with 0.8 power ranged from ~ 80 to >1000 samples (Fig. 4a) and ~ 40 to 60 samples for the CRSZ (Fig. 4b). Teleost N_{max} model outputs revealed the majority of years (2 of 4 at WHSZ; 2 of 3 at CRSZ) had low power (<0.7) to detect 50% change, independent of sample size. At the WHSZ, 2014 data were modelled to achieve >0.8

power with ~150 samples (~ 75 within project and reference), and 2013 data would achieve this level with ~1700 samples (~ 850 within project and reference) (Fig. 4c). The only data to achieve 0.8 power to detect 50% change in N_{max} at CRSZ was for 2012, which would require ~200 samples (100 within project and reference) (Fig. 4d).

Assemblage, species richness and abundance analyses

Comparison between study zones

Assemblage composition and species richness (S) models revealed significant differences between study zones ($P(\text{perm}) = <0.001$; Table 2). The species contributing most (top 4 species in SIMPER table) to the dissimilarity in composition were Poor cod *Trisopterus minutus*, common hermit crab *Pagurus bernhardus*, spiny starfish *Marthasterias glacialis* and cuckoo wrasse *Labrus mixtus*, with higher abundance of all except hermit crab in the CRSZ (Table 2). Higher average species richness was also observed within the CRSZ (CRSZ = mean 6.8 ± 2.2 SD, $n=107$; WHSZ = mean 5.3 ± 2.1 SD, $n=91$). Yearly variation in overall species richness and associated variance was observed in both study zones (Fig. 5), but was particularly evident in teleost abundance data (Fig. 6).

Table 2. PERMANOVA results for species assemblage and richness (S) models by study zones.
Fixed effects are non-italicized and random effects italicized. Significant permutation p-values below the 0.01 level for fixed effects are shown in bold.

Wave Hub vs Cable Route Zones							
<i>Assemblage</i>					<i>S</i>		
Source	d.f.	MS	Pseudo- <i>F</i>	<i>P</i> (perm)	MS	Pseudo- <i>F</i>	<i>P</i> (perm)
Zone	1	67326	13.159	0.0001	2.6850	8.0528	0.0029
<i>Treatment</i>	4	10121	3.2071	0.0017	1.0221	6.6190	0.0095
<i>Season</i>	1	4128.7	2.6529	0.0060	0.1888	2.0100	0.1602
<i>Location</i>	1	1144.5	0.3634	0.9417	0.2575	1.1326	0.3050
<i>Location(Treatment)</i>	21	3318.0	2.3059	0.0001	0.1961	2.2496	0.0019
Pooled terms	169	1439.0			0.0871		
Pair-wise test for Zone:					Pair-wise test for Zone:		
Group		t		<i>P</i> (perm)	Group	t	<i>P</i> (perm)
WH vs CR		3.6276		0.0001	WH vs CR	2.8378	0.0040
SIMPER output for Zone (>5% contribution):							
Species/family		WH av. abundance	CR av. abundance	% cont.			
<i>Trisopterus minutus</i>		1.25	1.44	11.4			
<i>Pagurus bernhardus</i>		1.18	0.16	8.47			
<i>Marthasterias glacialis</i>		0.28	1.18	8.23			
<i>Labrus mixtus</i>		0.25	1.24	8.07			
<i>Ophiurodea</i>		0.91	0.33	7.48			
<i>Ctenolabrus rupestris</i>		0.02	0.85	6.25			
<i>Scyliorhinus canicula</i>		1.08	0.91	5.87			
<i>Gobiidae</i>		0.52	0.43	5.05			

Fig. 5. Species richness (mean \pm standard error) and variance for each year at the (a) Wave Hub and (b) Cable Route study zones. Number above bar = sample size.





Fig. 6. Teleost relative abundance (mean \pm standard error) and variance for each year at the (a) Wave Hub and (b) Cable Route study zones. Number above bar = sample size.



Within Wave Hub study zone

No significant fixed effect or interaction term was found in the multivariate assemblage composition models for all species (all $P(\text{perm})$ values >0.05 ; Table 3a; Fig. S5), teleosts (all $P(\text{perm})$ values >0.05 ; Table S2a; Fig. S6) or crustaceans (all $P(\text{perm})$ values >0.05 ; Table S3a; Fig. S6) for the WHSZ.

Species richness (S) had a significant interaction $\text{Year} \times \text{Treatment}$ term ($P(\text{perm}) = 0.0136$; Table 3a), as did teleosts ($P(\text{perm}) = 0.0367$; Table S2a) and crustacean models ($P(\text{perm}) = 0.001$; Table S3a). Pairwise tests revealed that these were mainly driven by significant differences between years within the Wave Hub project and reference locations (Tables 3a, S2a & S3a), with only a significant effect of *Treatment* for all species in 2012 ($P = <0.05$; Table 3a) and for crustaceans in 2014 ($P = <0.001$; Table S3a). Relative abundance (N_{max}) models had no significant fixed effect or interaction term for teleosts, crustaceans,

echinoderms, elasmobranchs or *Pollachius spp.* (all $P(\text{perm})$ values = >0.05 ; Tables 3a, S2a, S3a, S5a & S5a), but there was a significant interaction term ($\text{Year} \times \text{Treatment}$) for *Cancer pagurus* ($P(\text{perm}) = <0.05$; Table S5a). Pairwise tests showed significant differences between years within Wave Hub project (2011-2013 & 2012-2013) or reference locations (2011-2012), and for locations within one year (2011), however some pairwise tests could not be conducted indicating a limitation of the data.

Within Cable Route study zone

Multivariate assemblage composition analysis revealed a significant interaction term, $\text{Year} \times \text{Treatment}$, for all species ($P(\text{perm}) = 0.0421$; Table 3b; Fig. S5) and teleost models ($P(\text{perm}) = 0.0118$; Table S2b; Fig. S6). Pairwise tests revealed both had significant differences between years within Cable Route project and reference locations, but not between Cable Route project and reference locations within years (Table 3b & Table S2b). The crustacean model had a significant Year term ($P(\text{perm}) = 0.0137$; Table S3b; Fig. S6), and 3 out of the 6 subsequent pairwise tests were significant (2011-13, 2011-14 & 2013-14; Table S3b). Species richness (S) models had no significant fixed terms for all species or teleosts (Table 3b & Table S2b), but the crustacean model revealed Year as significant ($P = <0.001$; Table S3b). Four out of the 6 subsequent pairwise tests were significant (2011-13, 2012-13, 2012-14 & 2013-14; Table S3b). Relative abundance (N_{max}) models had no significant fixed terms for teleosts, echinoderms or elasmobranchs (all $P(\text{perm})$ values >0.05 ; Table S2b and S4b). Year was found to be significant in the crustacean ($P(\text{perm}) = 0.0002$; Table S3a) and *Cancer pagurus* ($P(\text{perm}) = 0.0002$; Table S5b) models, with 4 from 6 pairwise test significant for crustaceans (2011-14, 2012-13, 2012-14, 2013-14; Table S3a) and 3 from 6 pairwise test significant for *Cancer pagurus* (2011-13, 2012-13, 2013-14; Table S5b). The only model to have a significant fixed Treatment term was for relative abundance of *Pollachius spp.*

454 ($P(\text{perm}) = 0.01$; Table S5b), with significantly greater abundance in Cable Route project
455 than reference locations (Cable Route: mean 1.4 ± 0.8 SD, $n=23$; reference = mean 1 ± 0 SD,
456 $n=7$).

457

458 Habitats

459 Assemblage composition and species richness (S) models revealed significant differences
460 between habitat types (Types 1-3: $P(\text{perm}) = <0.001$; Table 4 ; Fig. S7.). No significant
461 difference in N_{\max} was found between habitats.

462 **Table 3.** PERMANOVA results for all species assemblage, richness (S) and teleost species relative abundance (N_{max}) models for the Wave hub (a) and Cable
463 Route (b) study zones. Fixed effects are non-italicized and random effects italicized. Significant permutation p-values below the 0.05 level for fixed effects
464 and interactions are shown in bold.

(a) Wave Hub study zone										
<i>Assemblage (all species)</i>					<i>S (all species)</i>			<i>N_{max} (teleost species)</i>		
Source	d.f.	MS	Pseudo- <i>F</i>	<i>P</i> (perm)	MS	Pseudo- <i>F</i>	<i>P</i> (perm)	MS	Pseudo- <i>F</i>	<i>P</i> (perm)
Year	4	9525.1	2.6593	0.0838	1.26910	18.190	0.1155	4.6522	16.404	0.1225
Treatment	1	3424.5	0.7086	0.6604	0.00532	0.1248	0.7077	2.2460	1.5607	0.2674
<i>Season</i>	<i>1</i>	<i>3532.5</i>	<i>2.2758</i>	0.0207	<i>0.06245</i>	<i>0.7359</i>	<i>0.4073</i>	<i>0.2483</i>	<i>0.2947</i>	<i>0.5889</i>
<i>Habitat</i>	<i>3</i>	<i>5229.8</i>	<i>2.9543</i>	0.0001	<i>0.54909</i>	<i>6.9326</i>	0.0007	<i>1.8946</i>	<i>2.2091</i>	<i>0.0929</i>
<i>Location(Treatment)</i>	<i>6</i>	<i>4125.6</i>	<i>3.0025</i>	0.0001	<i>0.14502</i>	<i>2.1285</i>	<i>0.0588</i>	<i>1.2010</i>	<i>1.5015</i>	<i>0.1880</i>
Year x Treatment	4	1930.3	1.4048	0.0759	0.23181	3.4023	0.0136	1.2459	1.5576	0.1913
Pooled terms	71	1374.1			0.06813			0.7999		
Pair-wise test for Year x Treatment:										
<u>Project</u>										
Year					t		<i>P</i> (perm)			
2011-2012					2.7615		0.0155			
2011-2013					1.6244		0.1254			
2011-2014					3.9172		0.0375			
2011-2015					2.4403		0.0292			
2012-2013					10.346		0.0380			
2012-2014					2.7358		0.0278			
2012-2015					0.6682		0.5179			
2013-2014					7.1531		0.0003			
2013-2015					15.652		0.0481			
2014-2015					1.8411		0.0999			
<u>Reference</u>										
2011-2012					0.7015		0.5037			
2011-2013					0.9376		0.3630			
2011-2014					5.2210		0.0003			
2011-2015					2.5880		0.0211			
2012-2013					1.2550		0.4944			
2012-2014					5.4939		0.0002			
2012-2015					3.5548		0.0019			
2013-2014					3.7698		0.0489			
2013-2015					1.2859		0.4656			
2014-2015					3.7541		0.0046			
Pair-wise test for Year x Treatment :										
Year					t		<i>P</i> (perm)			
2011					1.4630		0.2122			

	2012	2.9688	0.0181	
	2013	1.5870	0.2158	
	2014	0.6848	0.5935	
	2015	0.5256	0.6315	

(b)

		Cable Route zone								
		<i>Assemblage (all species)</i>			<i>S (all species)</i>			<i>N_{max} (teleost species)</i>		
Source	d.f.	MS	Pseudo- <i>F</i>	<i>P</i> (perm)	MS	Pseudo- <i>F</i>	<i>P</i> (perm)	MS	Pseudo- <i>F</i>	<i>P</i> (perm)
Year	3	6536.5	2.5807	0.0079	0.33179	1.0787	0.4573	3.7770	1.1392	0.4471
Treatment	1	2513.6	1.1766	0.3133	0.38619	1.7755	0.2085	2.6261	2.5411	0.1362
Season	1	3245.0	2.0852	0.0369	0.46417	6.6699	0.0126	5.3069	7.8565	0.0060
Zone	1	3532.7	1.8065	0.0743	0.04206	0.2403	0.6855	0.4412	0.5121	0.5018
Habitat	3	3532.2	2.6270	0.0002	0.07837	0.9400	0.4225	1.4894	2.3797	0.0794
Location(Treatment)	13	1823.3	1.4602	0.0038	0.18391	2.9089	0.0027	0.8614	1.4885	0.1439
Year x Treatment	3	1953.6	1.5646	0.0421	0.10769	1.7034	0.1742	1.1476	1.9832	0.1243
Pooled terms	81	1248.7			0.06322			0.5787		
Pair-wise test for Year x Treatment:										
<u>Project</u>										
Year			t	<i>P</i> (perm)						
2011-2012			2.1595	0.0001						
2011-2013			1.4024	0.0473						
2011-2014			1.8228	0.0046						
2012-2013			2.9543	0.0007						
2012-2014			1.2262	0.2050						
2013-2014			1.8894	0.0062						
<u>Reference</u>										
2011-2012			1.2666	0.1783						
2011-2013			1.2847	0.1559						
2011-2014			2.1578	0.0007						
2012-2013			0.8882	0.6195						
2012-2014			1.1486	0.2702						
2013-2014			1.4463	0.1088						
Pair-wise test for Year x Treatment :										
Year			t	<i>P</i> (perm)						
2011			0.9143	0.5935						
2012			0.8401	0.7480						
2013			1.2993	0.1492						
2014			0.8659	0.5963						

465

466

467 **Table 4.** PERMANOVA results for species assemblage, richness (S) and relative abundance (N_{max}) models by habitat types; 1-3. Fixed effects are non-
468 italicized and random effects italicized. Significant permutation p-values below the 0.01 level for fixed effects are shown in bold.

Habitats (1 – 3)										
<i>Assemblage</i>				<i>S</i>			<i>N_{max}</i>			
Source	d.f.	MS	Pseudo- <i>F</i>	<i>P</i> (perm)	MS	Pseudo- <i>F</i>	<i>P</i> (perm)	MS	Pseudo- <i>F</i>	<i>P</i> (perm)
Habitat	2	30502	15.549	0.0001	1.2972	11.763	0.0001	1.1099	1.8523	0.1561
<i>Year</i>	4	<i>8813.1</i>	<i>5.2586</i>	0.0001	<i>0.6820</i>	<i>7.8907</i>	0.0001	<i>4.2311</i>	<i>9.4497</i>	0.0001
Pooled terms	182	1675.9			0.0864			0.4477		
Pair-wise test for Habitat				Pair-wise test for Habitat						
Habitats		t	<i>P</i> (perm)	Habitats		t	<i>P</i> (perm)			
1-3		5.2528	0.0001	1-3		4.7223	0.0001			
1-2		2.5931	0.0001	1-2		2.5438	0.0123			
3-2		3.0235	0.0001	3-2		1.9184	0.0582			

Discussion

Weather and sea state are major considerations when working in mid latitude offshore marine environments, and can be a serious limitation to survey time and sampling effort when using small vessels. In this study, poor weather conditions and logistical constraints reduced the bi-annual survey to one season a year, with the exception of 2013. Access to larger vessels may have improved this situation but would have appreciably increased costs and reduced cost effectiveness of the BRUV technique. The study period encompassed some noteworthy bad weather events across the British Isles, and the extreme 2014 storms (Masselink et al., 2016) resulted in a reduction in samples due to bad visibility rendering footage unusable (Fig. S2). With storm events predicted to increase with climate change (Coumou and Rahmstorf, 2012; Zappa et al., 2013) the potential effects on sampling success need to be considered within survey design and sampling effort plans, i.e. over-estimate required sampling for planned redundancy. The coastal areas optimal for wind or wave energy generation will be, by their nature, open to weather systems that will influence access to locations (sampling opportunities) and the quality of data gathered (visibility). The sampling loss due to camera and battery failures was relatively high and likely related to the repeated demands placed on the equipment and experience of maintaining and deploying the equipment on/off small boats in challenging conditions with small research teams (2 people). These losses were minimised with experience and improved equipment, but could be further mitigated if funds and time are available for repeat sampling efforts and/or campaigns.

The loss of survey periods or samples due to weather, technical problems or logistical issues reduces the precision in the characterisation of spatial and temporal variability at a site, and will ultimately affect the ability to detect impact (Underwood and Chapman, 2003). Deploying BRUV systems for an optimal time can help reduce avoidable loss by limiting time in the water. Accumulation curves derived from data gathered at different locations

494 around the world, for fish and invertebrate species, suggests BRUV deployment time (or
495 footage time) ranging from 30 minutes to 2 hours in order to capture a significant proportion
496 of the number of species or individuals attracted to the bait (Bernard and Gotz, 2012;
497 Unsworth et al., 2014). BRUVs were deployed for 60 minutes in our study, but only the first
498 30 minutes of the footage were used in the analyses, capturing on average 75% or more of the
499 total richness or relative abundance observed over the whole 60 minute footage (Fig. 3). Our
500 decision is worth briefly elaborating as it considered ecological and statistical elements of
501 BRUV sampling and, therefore, could be informative for future studies using this technique.
502 Firstly, in our study a BRUV system was used to provide a rapid assessment of species and
503 relative abundance for particular, small benthic areas. If longer time periods were used, the
504 effective range of attraction (AR) would increase so mobile species (teleost and
505 elasmobranch) could be attracted from much further afield (>500 m; not our objective), and
506 as a consequence, the animals could potentially attend multiple cameras, leading to double
507 counting of individuals (pseudo-replication in our design). Secondly, due to technical
508 (camera/battery failure) and condition (visibility or field of view) related issues, the number
509 of useable videos significantly reduced (~40%) when applying 60 minute (116 videos)
510 compared to a 30 minute footage requirement (198 videos) (Fig. S1). The decrease in sample
511 size reduces the power of these data to detect statistically significant change and, for 2011,
512 would have translated into no data being available and removal of this year from analyses.
513 The statistical benefit conferred by an increased sample size and inclusion of all surveyed
514 years (inter-annual variance), combined with the ecological reasoning behind using a shorter
515 time period, resulted in the decision to use 30 minute duration video data. However, a
516 limitation of only deploying, or using data, for a short period (30 minutes in our case), is the
517 chance of missing the presence of rare and/or cautious species. Given renewable energy
518 development sites (wave, wind or tidal) could become *de facto* Marine Protection Areas

(MPAs) by excluding damaging seabed fishing activities, the areas could become refuges for species with small or recovering populations. Deploying a subset of BRUVs for longer could help detect such species and be incorporated into study designs, specifically for this purpose. In this study, no new species were detected when 60 minutes footage was compared to 30 minutes, but longer periods may be necessary to detect rare species. The specifics detailed here are particular to our study but highlight, more generally, the need to understand the survey technique being used and the data it is expected to capture. Moreover, how the precautionary approach to sampling effort previously recommended (over-estimation) is necessary to allow redundancy in sampling campaigns when working in highly dynamic environments.

The analyses conducted using the BRUV data provided little consistent evidence of differences in metrics (across taxa) between the locations influenced by either the trawling exclusion (Wave Hub) or cable infrastructure (Cable Route), and reference locations within study zones. The only consistent change across years was an increase in the relative abundance of pollack and saithe (*Pollachius spp.*) around the cable infrastructure. These are commercially important coastal species understood to associate with rocky reef and hard substrate habitat as nursery areas (Seitz et al., 2014), suggesting the addition of rock and concrete matressing on the cable may be providing suitable conditions. Greater abundance of cuckoo wrasse *Labrus mixtus* observed during towed camera surveys in the same area has also been attributed to the presence of the hard substrate for cable protection (Sheehan et al., 2013). Both observations are consistent with other studies of fish abundance around MREI structures (Wilhelmsson et al., 2006) and add to the evidence these introduced structures and associated infrastructure are created habitat for species to utilise (Inger et al., 2009; Miller et al., 2013; Sheehan et al., 2018; Witt et al., 2012). The lack of evidence towards an impact of trawling exclusion in the Wave Hub zone may relate to low fishing effort in the area before

(Campbell et al., 2013) and during the study, creating equivalent disturbance (or lack of) in project and reference locations. Alternatively, it could be an artefact of the survey design and power to detect change. The power analyses revealed how the ability to detect change in species richness and teleost relative abundance differed considerably between zones (spatial) and among survey years (temporal), producing a range of sampling schedules to detect the same effect size and varying levels of power (Fig. 4). Low power was particularly apparent in teleost abundance indicating these data could only confidently detect large changes (>50%) with high probability ($\alpha=0.05$), much less for the CRSZ. This could explain the lack of significant results found for this metric in our univariate analyses (Table 2 & Table S2), with potential influence on the outcome of the multivariate assemblage composition analysis (Fig. S5). Low probability to detect directional change in abundance data due to high variability has been found for other survey techniques and species (e.g. fish, cetaceans & seabirds; Al-Chokhachy et al., 2009; Forney, 2000; Maclean et al., 2013). Our results re-iterate caution in accepting that no change is taking place when analyses fail to reveal statistically significant patterns, when a lack of statistical power may be the contributing factor (Al-Chokhachy et al., 2009; Maclean et al., 2013). Despite good statistical power (~ 0.8) to detect relatively small changes (20%) in species richness, the majority of statistically significant effects found in our analyses (uni- and multivariate) were associated with yearly differences (Tables 2 and Tables S2-5), and only a single year difference found for the exclusion zone (*'Treatment'*) in richness of all species combined (2012; Table 2a) and crustaceans (2014; Table S3a) at the WHSZ. Natural yearly variation was the main effect identified at the study zones suggesting high levels of 'noise' in the system, presenting potential difficulty in distinguishing impacts related to the Wave Hub exclusion zone or cable route infrastructure. Moreover, it demonstrates how multi-year data are essential to capture site variability and provide accurate

baseline characterisation, from which single site or cumulative impact from renewable energy convertors or manufactured infrastructure could be robustly assessed (Maclean et al., 2014).

The high natural inter-annual variation found for each metric (at both study zones; Figs. 4-6) presents evidence for the difficulty in using pilot (e.g. single survey or one year) site data to inform the design and effort (e.g. power analysis) required for robust impact assessment surveys. If the power analysis was to rely on data from only one year of our study there would be a risk of either over- or under-estimating the number of samples needed to obtain acceptable power to detect a chosen effect size (Fig. 4). The consequence could either be a lack of power in the subsequent impact analyses (under-estimate) or collecting samples that are potentially unnecessary (over-estimate). The latter has the benefits of providing redundancy in the sampling effort, as has been previously recommended, but requires additional time, effort and cost, which would need to be factored into the assessment program.

The difference in assemblage composition, species richness and variance found between study zones (Table 2; Fig. 4 & 5) might be expected given the dominant habitat types found at the Wave Hub (medium to fine sediment) and Cable Route (mix of rocky reef and large boulders) study zones (e.g. Fig. S7), and the significant difference in composition and richness found among them (Table 4). The site specific nature of ecosystem components, even over a relatively small spatial area (~130 km²), is effectively demonstrated using these data. The characterisations these data offer also illustrate how useful consistent data collection methods (and survey design) can be in allowing comparison between locations. Baseline characterisation and impact studies of benthic habitats and species are required at MREIs located in coastal locations (inshore and offshore) with various physical (e.g. size, bathymetry) and ecological characteristics. While the energy convertor design may differ between locations the ability to understand more general or cumulative effects caused by

developments would be enhanced by multi-site data, collected using the same survey techniques. Adopting standard practices and guidelines for pre- and post-development benthic survey methods, design and analysis would help optimise costs associated with EIAs and, if adopted across multi-sites and multi-years, could ultimately lead to impact models with predictive power for species, communities or ecosystems (e.g. Butenschön et al., 2016).

Given there were no prior ecological data for the study zones monitored during this study, the survey design performed well in providing adequate samples to detect change in species richness but not so well for relative abundance. Caution is required when interpreting the abundance results (detailed above), but overall it indicates the survey would be able to detect differences between project and reference locations that could be considered large (e.g. >50%). Whether changes of this size are biologically or ecologically significant is unknown and would be highly dependent on the resilience of the ecosystem. For example, if a reduction in species richness included the loss of a functional group (e.g. predators or herbivores) within the local ecosystem, then it is likely this could lead to a significant ecological impact (Micheli and Halpern, 2005).

As part of this study the BRUV system demonstrated its value as a tool for collecting assemblage composition, species richness and relative abundance data for epi-benthic mobile species in highly dynamic conditions, and a good candidate for use as part of marine EIAs across latitudes. The system offers a cost effective and flexible method that can provide the spatial and temporal coverage that is difficult to obtain using other methods (i.e. divers, remote underwater vehicles). When used with stereo cameras, BRUV can also offer size data that could help elucidate more detailed age related effects cause by introduced or altered habitat (Elliott et al., 2017), or converted into biomass estimates (standing or relative) providing another metric to assess impact. Although other traditional methods of sampling these communities, e.g. trawling, potting, bottom lines or nets, can provide these metrics and

work in similar or worse sea conditions, they can be destructive or taxa specific, increasing the cost of survey effort and to the ecosystem of study. With further development, BRUV systems also have the potential to help address data collection gaps surveys often suffer from, e.g. diurnal variation. With the integration of movement sensors (i.e. infra-red) and/or artificial intelligent (AI) algorithms to activate equipment on species presence, 24 hour deployments could be possible. BRUV biodiversity data has also been found to be complimentary to environmental DNA (Stat et al., 2018), suggesting a combined approach using these non-invasive methods could further enhance the effectiveness of monitoring surveys.

Acknowledgments

This work was funded by a Natural Environment Research Grant (NE/J012319/1), an Engineering and Physical Sciences Research Council Grant (EP/P026109/1), an Intelligent Energy Europe (IEE) Grant (IEE/09/809/SI2.558291) and a European Cross-border Cooperation INTERREG IV Grant, co-founded by the European Regional Development Fund (ERDF). Thank you to Chris Lowe (Newquay Sea Safaris and Fishing) and Lech Kwiatkowski for safely skippering survey vessels, and two anonymous reviewers for their helpful comments and suggestions.

References

- Al-Chokhachy, R., Budy, P., Conner, M., 2009. Detecting declines in the abundance of a bull trout (*Salvelinus confluentus*) population: understanding the accuracy, precision, and costs of our efforts. *Canadian Journal of Fisheries and Aquatic Sciences* 66, 649-658.
- Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecology* 26, 32-46.

643 Anderson, M.J., Millar, R.B., 2004. Spatial variation and effects of habitat on temperate reef
644 fish assemblages in northeastern New Zealand. *Journal of Experimental Marine Biology and*
645 *Ecology* 305, 191-221.

646 Bailey, H., Brookes, K.L., Thompson, P.M., 2014. Assessing environmental impacts of
647 offshore wind farms: lessons learned and recommendations for the future. *Aquatic*
648 *biosystems* 10, 8.

649 Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models
650 Using lme4. *Journal of Statistical Software* 67, 1-48.

651 Bernard, A.T.F., Gotz, A., 2012. Bait increases the precision in count data from remote
652 underwater video for most subtidal reef fish in the warm-temperate Agulhas bioregion.
653 *Marine Ecology Progress Series* 471, 235-252.

654 Bicknell, A.W.J., Godley, B.J., Sheehan, E.V., Votier, S.C., Witt, M.J., 2016. Camera
655 technology for monitoring marine biodiversity and human impact. *Frontiers in Ecology and*
656 *the Environment* 14, 424-432.

657 Boehlert, G.W., Gill, A.B., 2010. Environmental and ecological effects of ocean renewable
658 energy development: a current synthesis. *Oceanography* 23, 68-81.

659 Bray, J.R., Curtis, J.T., 1957. An ordination of the upland forest communities of southern
660 Wisconsin. *Ecological monographs* 27, 325-349.

661 Bulleri, F., Chapman, M.G., 2010. The introduction of coastal infrastructure as a driver of
662 change in marine environments. *Journal of Applied Ecology* 47, 26-35.

663 Butenschön, M., Clark, J., Aldridge, J.N., Allen, J.I., Artioli, Y., Blackford, J., Bruggeman,
664 J., Cazenave, P., Ciavatta, S., Kay, S., Lessin, G., van Leeuwen, S., van der Molen, J., de
665 Mora, L., Polimene, L., Sailley, S., Stephens, N., Torres, R., 2016. ERSEM 15.06: a generic
666 model for marine biogeochemistry and the ecosystem dynamics of the lower trophic levels.
667 *Geosci. Model Dev.* 9, 1293-1339.

668 Campbell, M.S., Stehfest, K.M., Votier, S.C., Hall-Spencer, J.M., 2013. Mapping fisheries
669 for marine spatial planning: Gear-specific vessel monitoring system (VMS), marine
670 conservation and offshore renewable energy. *Marine Policy*.

671 Cappo, M., Harvey, E., Malcolm, H., Speare, P., 2003. Potential of video techniques to
672 monitor diversity, abundance and size of fish in studies of marine protected areas, in:
673 Beumer, J.P., Grant, A., Smith, D.C. (Eds.), *Aquatic Protected Areas-what works best and*
674 *how do we know?*, Cairns, Queensland, Australia, pp. 455-464.

675 Cappo, M., Speare, P., De'ath, G., 2004. Comparison of baited remote underwater video
676 stations (BRUVS) and prawn (shrimp) trawls for assessments of fish biodiversity in inter-
677 reefal areas of the Great Barrier Reef Marine Park. *Journal of Experimental Marine Biology*
678 *and Ecology* 302, 123-152.

679 Clarke, K., Chapman, M., Somerfield, P., Needham, H., 2006a. Dispersion-based weighting
680 of species counts in assemblage analyses. *Marine Ecology Progress Series* 320, 11-27.

681 Clarke, K.R., Somerfield, P.J., Chapman, M.G., 2006b. On resemblance measures for
682 ecological studies, including taxonomic dissimilarities and a zero-adjusted Bray–Curtis
683 coefficient for denuded assemblages. *Journal of Experimental Marine Biology and Ecology*
684 330, 55-80.

685 Clarke, K.R., Warwick, R.M., 2001. *Change in marine communities: an approach to*
686 *statistical analysis and interpretation*. 2nd Edition. PRIMER-E Ltd: Plymouth, United
687 Kingdom.

688 Costanza, R., d'Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K.,
689 Naeem, S., O'Neill, R.V., Paruelo, J., Raskin, R.G., Sutton, P., van den Belt, M., 1997. The
690 value of the world's ecosystem services and natural capital. *Nature* 387, 253.

691 Coumou, D., Rahmstorf, S., 2012. A decade of weather extremes. *Nature Climate Change* 2,
692 491.

693 Denny, C.M., Willis, T.J., Babcock, R.C., 2004. Rapid recolonisation of snapper *Pagrus*
694 *auratus*: Sparidae within an offshore island marine reserve after implementation of no-take
695 status. *Marine Ecology Progress Series* 272, 183-190.

696 Dorman, S.R., Harvey, E.S., Newman, S.J., 2012. Bait Effects in Sampling Coral Reef Fish
697 Assemblages with Stereo-BRUVs. *PLoS ONE* 7, e41538.

698 Drakou, E.G., Pendleton, L., Effron, M., Ingram, J.C., Teneva, L., 2017. When ecosystems
699 and their services are not co-located: oceans and coasts. *ICES Journal of Marine Science* 74,
700 1531-1539.

701 Eigaard, O.R., Bastardie, F., Breen, M., Dinesen, G.E., Hintzen, N.T., Laffargue, P.,
702 Mortensen, L.O., Nielsen, J.R., Nilsson, H.C., O'Neill, F.G., Polet, H., Reid, D.G., Sala, A.,
703 Sköld, M., Smith, C., Sørensen, T.K., Tully, O., Zengin, M., Rijnsdorp, A.D., 2015.
704 Estimating seabed pressure from demersal trawls, seines, and dredges based on gear design
705 and dimensions. *ICES Journal of Marine Science: Journal du Conseil*.

706 Elliott, S.A.M., Turrell, W.R., Heath, M.R., Bailey, D.M., 2017. Juvenile gadoid habitat and
707 ontogenetic shift observations using stereo-video baited cameras. *Marine Ecology Progress*
708 *Series* 568, 123-135.

709 Ellis, D.M., DeMartini, E.E., 1995. Evaluation of a video camera technique for indexing
710 abundances of juvenile pink snapper, *Pristipomoides filamentosus*, and other Hawaiian
711 insular shelf fishes. *Oceanographic Literature Review* 42.

712 Forney, K.A., 2000. Environmental Models of Cetacean Abundance: Reducing Uncertainty
713 in Population Trends. *Conservation Biology* 14, 1271-1286.

714 Fox, C.J., Benjamins, S., Masden, E.A., Miller, R., 2018. Challenges and opportunities in
715 monitoring the impacts of tidal-stream energy devices on marine vertebrates. *Renewable and*
716 *Sustainable Energy Reviews* 81, 1926-1938.

717 Franco, A., Quintino, V., Elliott, M., 2015. Benthic monitoring and sampling design and
 718 effort to detect spatial changes: A case study using data from offshore wind farm sites.
 719 Ecological Indicators 57, 298-304.

720 Galparsoro, I., Borja, A., Uyarra, M.C., 2014. Mapping ecosystem services provided by
 721 benthic habitats in the European North Atlantic Ocean. Frontiers in Marine Science 1.

722 Gill, A.B., 2005. Offshore renewable energy: ecological implications of generating electricity
 723 in the coastal zone. Journal of Applied Ecology 42, 605-615.

724 Green, P., MacLeod, C.J., 2016. SIMR: an R package for power analysis of generalized linear
 725 mixed models by simulation. Methods in Ecology and Evolution 7, 493-498.

726 Griffin, R.A., Robinson, G.J., West, A., Gloyne-Phillips, I.T., Unsworth, R.K.F., 2016.
 727 Assessing Fish and Motile Fauna around Offshore Windfarms Using Stereo Baited Video.
 728 PLoS ONE 11, e0149701.

729 GWEC, 2016. Global Wind Report: Annual Market Update 2015, p. 76.

730 Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agrosa, C., Bruno,
 731 J.F., Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin,
 732 E.M.P., Perry, M.T., Selig, E.R., Spalding, M., Steneck, R., Watson, R., 2008. A Global Map
 733 of Human Impact on Marine Ecosystems. Science 319, 948-952.

734 Harvey, E., Santana Garcon, J., Goetze, J., Saunders, B., Cappel, M., 2018. The Use of
 735 Stationary Underwater Video for Sampling Sharks, in: Carrier, J., Heithaus, M.R.,
 736 Simpfendorfer, C.A. (Eds.), Shark Research: Emerging Technologies and Applications for
 737 the Field and Laboratory. CRC Press, pp. 111-132.

738 Harvey, E.S., Butler, J.J., McLean, D.L., Shand, J., 2012. Contrasting habitat use of diurnal
 739 and nocturnal fish assemblages in temperate Western Australia. Journal of Experimental
 740 Marine Biology and Ecology 426-427, 78-86.

741 Harvey, E.S., Cappo, M., Butler, J.J., Hall, N., Kendrick, G.A., 2007. Bait attraction affects
 742 the performance of remote underwater video stations in assessment of demersal fish
 743 community structure. *Marine Ecology Progress Series* 350, 245-254.

744 Heagney, E.C., Lynch, T.P., Babcock, R.C., Suthers, I.M., 2007. Pelagic fish assemblages
 745 assessed using mid-water baited video: standardising fish counts using bait plume size.
 746 *Marine Ecology Progress Series* 350, 255.

747 Heery, E.C., Bishop, M.J., Critchley, L.P., Bugnot, A.B., Airoidi, L., Mayer-Pinto, M.,
 748 Sheehan, E.V., Coleman, R.A., Loke, L.H.L., Johnston, E.L., Komyakova, V., Morris, R.L.,
 749 Strain, E.M.A., Naylor, L.A., Dafforn, K.A., 2017. Identifying the consequences of ocean
 750 sprawl for sedimentary habitats. *Journal of Experimental Marine Biology and Ecology* 492,
 751 31-48.

752 Hiddink, J.G., Jennings, S., Sciberras, M., Szostek, C.L., Hughes, K.M., Ellis, N., Rijnsdorp,
 753 A.D., McConnaughey, R.A., Mazon, T., Hilborn, R., Collie, J.S., Pitcher, C.R., Amoroso,
 754 R.O., Parma, A.M., Suuronen, P., Kaiser, M.J., 2017. Global analysis of depletion and
 755 recovery of seabed biota after bottom trawling disturbance. *Proceedings of the National*
 756 *Academy of Sciences*.

757 Inger, R., Attrill, M.J., Bearhop, S., Broderick, A.C., James Grecian, W., Hodgson, D.J.,
 758 Mills, C., Sheehan, E., Votier, S.C., Witt, M.J., Godley, B.J., 2009. Marine renewable energy:
 759 potential benefits to biodiversity? An urgent call for research. *Journal of Applied Ecology* 46,
 760 1145-1153.

761 Jackson, E.L., Langmead, O., Barnes, M., Tyler-Walters, H., Hiscock, K., 2009.
 762 Identification of indicator species to represent the full range of benthic life history strategies
 763 for Lyme Bay and the consideration of the wider application for monitoring of Marine
 764 Protected Areas.

765 Judd, A., 2012. Guidelines for data acquisition to support marine environmental assessments
 766 of offshore renewable energy projects. CEFAS, Lowestoft.

767 Langhamer, O., Wilhelmsson, D., 2009. Colonisation of fish and crabs of wave energy
 768 foundations and the effects of manufactured holes—a field experiment. *Marine Environmental*
 769 *Research* 68, 151-157.

770 Liqueste, C., Piroddi, C., Drakou, E.G., Gurney, L., Katsanevakis, S., Charef, A., Egoh, B.,
 771 2013. Current Status and Future Prospects for the Assessment of Marine and Coastal
 772 Ecosystem Services: A Systematic Review. *PLOS ONE* 8, e67737.

773 Lowry, M., Folpp, H., Gregson, M., Suthers, I., 2012. Comparison of baited remote
 774 underwater video (BRUV) and underwater visual census (UVC) for assessment of artificial
 775 reefs in estuaries. *Journal of Experimental Marine Biology and Ecology* 416–417, 243-253.

776 Maclean, I.M.D., Inger, R., Benson, D., Booth, C.G., Embling, C.B., Grecian, W.J.,
 777 Heymans, J.J., Plummer, K.E., Shackshaft, M., Sparling, C.E., Wilson, B., Wright, L.J.,
 778 Bradbury, G., Christen, N., Godley, B.J., Jackson, A.C., McCluskie, A., Nicholls-Lee, R.,
 779 Bearhop, S., 2014. Resolving issues with environmental impact assessment of marine
 780 renewable energy installations. *Frontiers in Marine Science* 1.

781 Maclean, I.M.D., Rehfisch, M.M., Skov, H., Thaxter, C.B., 2013. Evaluating the statistical
 782 power of detecting changes in the abundance of seabirds at sea. *Ibis* 155, 113-126.

783 Malcolm, H.A., Gladstone, W., Lindfield, S., Wraith, J., Lynch, T.P., 2007. Spatial and
 784 temporal variation in reef fish assemblages of marine parks in New South Wales, Australia-
 785 baited video observations. *Marine Ecology Progress Series* 350, 277.

786 Mallet, D., Pelletier, D., 2014. Underwater video techniques for observing coastal marine
 787 biodiversity: A review of sixty years of publications (1952–2012). *Fisheries Research* 154,
 788 44-62.

789 Masselink, G., Castelle, B., Scott, T., Dodet, G., Suanez, S., Jackson, D., Floc'h, F., 2016.
 790 Extreme wave activity during 2013/2014 winter and morphological impacts along the
 791 Atlantic coast of Europe. *Geophysical Research Letters* 43, 2135-2143.
 792 Micheli, F., Halpern, B.S., 2005. Low functional redundancy in coastal marine assemblages.
 793 *Ecology Letters* 8, 391-400.
 794 Millennium Ecosystem Assessment, 2005. *Ecosystems and Human Well-Being: Synthesis*.
 795 World Resources Institute, Washington, DC.
 796 Miller, R.G., Hutchison, Z.L., Macleod, A.K., Burrows, M.T., Cook, E.J., Last, K.S., Wilson,
 797 B., 2013. Marine renewable energy development: assessing the Benthic Footprint at multiple
 798 scales. *Frontiers in Ecology and the Environment* 11, 433-440.
 799 Neumann, B., Vafeidis, A.T., Zimmermann, J., Nicholls, R.J., 2015. Future Coastal
 800 Population Growth and Exposure to Sea-Level Rise and Coastal Flooding - A Global
 801 Assessment. *PLoS ONE* 10, e0118571.
 802 Osenberg, C.W., Schmitt, R.J., Holbrook, S.J., Abu-Saba, K.E., Flegal, A.R., 1994. Detection
 803 of Environmental Impacts: Natural Variability, Effect Size, and Power Analysis. *Ecological*
 804 *Applications* 4, 16-30.
 805 Pearce, B., Fariñas-Franco, J.M., Wilson, C., Pitts, J., deBurgh, A., Somerfield, P.J., 2014.
 806 Repeated mapping of reefs constructed by *Sabellaria spinulosa* Leuckart 1849 at an offshore
 807 wind farm site. *Continental Shelf Research* 83, 3-13.
 808 Priede, I.G., Bagley, P.M., Smith, A., Creasey, S., Merrett, N.R., 1994. Scavenging deep
 809 demersal fishes of the Porcupine Seabight, North-east Atlantic: observations by baited
 810 camera, trap and trawl. *Journal of the Marine Biological Association of the United Kingdom*
 811 74, 481-498.
 812 R Development Core Team, 2011. *R: A Language and Environment for Statistical*
 813 *Computing*. R Foundation for Statistical Computing, Vienna, Austria.

814 Seitz, R.D., Wennhage, H., Bergström, U., Lipcius, R.N., Ysebaert, T., 2014. Ecological
815 value of coastal habitats for commercially and ecologically important species. *ICES Journal*
816 *of Marine Science* 71, 648-665.

817 Sheehan, E.V., Cartwright, A.Y., Witt, M.J., Attrill, M.J., Vural, M., Holmes, L.A., Handling
818 editor: Silvana, B., 2018. Development of epibenthic assemblages on artificial habitat
819 associated with marine renewable infrastructure. *ICES Journal of Marine Science*, fsy151-
820 fsy151.

821 Sheehan, E.V., Stevens, T.F., Gall, S.C., Cousens, S.L., Attrill, M.J., 2014. Recovery of a
822 Temperate Reef Assemblage in a Marine Protected Area following the Exclusion of Towed
823 Demersal Fishing. *PLOS ONE* 8, e83883.

824 Sheehan, E.V., Witt, M.J., Cousens, S.L., Gall, S.C., Attrill, M.J., 2013. Benthic Interactions
825 with Renewable Energy Installations in a Temperate Ecosystem, The Twenty-third
826 International Offshore and Polar Engineering Conference. International Society of Offshore
827 and Polar Engineers.

828 Stat, M., John, J., DiBattista, J.D., Newman, S.J., Bunce, M., Harvey, E.S., 2018. Combined
829 use of eDNA metabarcoding and video surveillance for the assessment of fish biodiversity.
830 *Conservation Biology* 0.

831 Stenberg, C., Støttrup, J.G., van Deurs, M., Berg, C.W., Dinesen, G.E., Mosegaard, H.,
832 Grome, T.M., Leonhard, S.B., 2015. Long-term effects of an offshore wind farm in the North
833 Sea on fish communities. *Marine Ecology Progress Series* 528, 257-265.

834 Stobart, B., Díaz, D., Álvarez, F., Alonso, C., Mallol, S., Goñi, R., 2015. Performance of
835 baited underwater video: does it underestimate abundance at high population densities? *PloS*
836 *one* 10, e0127559.

837 Stobart, B., García-Charton, J.A., Espejo, C., Rochel, E., Goñi, R., Reñones, O., Herrero, A.,
838 Crec'hriou, R., Polti, S., Marcos, C., Planes, S., Pérez-Ruzafa, A., 2007. A baited underwater

839 video technique to assess shallow-water Mediterranean fish assemblages: Methodological
840 evaluation. *Journal of Experimental Marine Biology and Ecology* 345, 158-174.

841 Underwood, A.J., Chapman, M.G., 2003. Power, precaution, Type II error and sampling
842 design in assessment of environmental impacts. *Journal of Experimental Marine Biology and*
843 *Ecology* 296, 49-70.

844 Unsworth, R.K.F., Peters, J.R., McCloskey, R.M., Hinder, S.L., 2014. Optimising stereo
845 baited underwater video for sampling fish and invertebrates in temperate coastal habitats.
846 *Estuarine, Coastal and Shelf Science* 150, Part B, 281-287.

847 Ward, R.C., Loftis, J.C., McBride, G.B., 1986. The “data-rich but information-poor”
848 syndrome in water quality monitoring. *Environmental Management* 10, 291-297.

849 Watson, D.L., Anderson, M.J., Kendrick, G.A., Nardi, K., Harvey, E.S., 2009. Effects of
850 protection from fishing on the lengths of targeted and non-targeted fish species at the
851 Houtman Abrolhos Islands, Western Australia. *Marine Ecology Progress Series* 384, 241-
852 249.

853 Watson, D.L., Harvey, E.S., Kendrick, G.A., Nardi, K., Anderson, M.J., 2007. Protection
854 from fishing alters the species composition of fish assemblages in a temperate-tropical
855 transition zone. *Marine Biology* 152, 1197-1206.

856 Whitmarsh, S.K., Fairweather, P.G., Huvaneers, C., 2016. What is Big BRUVver up to?
857 Methods and uses of baited underwater video. *Reviews in Fish Biology and Fisheries*, 1-21.

858 Wilding, T.A., Gill, A.B., Boon, A., Sheehan, E., Dauvin, J.C., Pezy, J.-P., O’Beirn, F.,
859 Janas, U., Rostin, L., De Mesel, I., 2017. Turning off the DRIP (‘Data-rich, information-
860 poor’) – rationalising monitoring with a focus on marine renewable energy developments and
861 the benthos. *Renewable and Sustainable Energy Reviews* 74, 848-859.

862 Wilhelmsson, D., Malm, T., Öhman, M.C., 2006. The influence of offshore windpower on
863 demersal fish. *ICES Journal of Marine Science: Journal du Conseil* 63, 775-784.

864 Willis, T.J., Babcock, R.C., 2000. A baited underwater video system for the determination of
865 relative density of carnivorous reef fish. *Marine and Freshwater Research* 51, 755-763.

866 WindEurope, 2017. The European offshore wind industry: Key trends and statistics 2016, p.
867 37.

868 Witt, M.J., Sheehan, E.V., Bearhop, S., Broderick, A.C., Conley, D.C., Cotterell, S.P., Crow,
869 E., Grecian, W.J., Halsband, C., Hodgson, D.J., Hosegood, P., Inger, R., Miller, P.I., Sims,
870 D.W., Thompson, R.C., Vanstaen, K., Votier, S.C., Attrill, M.J., Godley, B.J., 2012.
871 Assessing wave energy effects on biodiversity: the Wave Hub experience. *Philosophical*
872 *Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 370,
873 502-529.

874 Zappa, G., Shaffrey, L.C., Hodges, K.I., Sansom, P.G., Stephenson, D.B., 2013. A
875 Multimodel Assessment of Future Projections of North Atlantic and European Extratropical
876 Cyclones in the CMIP5 Climate Models. *Journal of Climate* 26, 5846-5862.

877 Zintzen, V., Anderson, M.J., Roberts, C.D., Harvey, E.S., Stewart, A.L., Struthers, C.D.,
878 2012. Diversity and Composition of Demersal Fishes along a Depth Gradient Assessed by
879 Baited Remote Underwater Stereo-Video. *PLoS ONE* 7, e48522.